



Pregnancy Loss and Infant Death Alliance

SUPPORT . ADVOCACY . AWARENESS . EDUCATION

Position Statement

**Offer the Baby to Bereaved Parents With
Relationship-Based Care**

Position

Offer the Baby to Bereaved Parents With Relationship-Based Care

Offering the baby to bereaved parents should occur within the context of relationship-based care, supportive conversations with parents, and respecting their autonomy.

Executive Summary

When a baby dies, parents should be offered their child within the context of an empathic relationship, where the healthcare provider engages parents in periodic conversations, eliciting their thoughts and feelings about spending time with their baby. This engagement should include supporting the parents in doing what they believe is best for their family, whether or not they elect to spend time with their baby.

- When a baby dies during pregnancy, birth, or following delivery, healthcare providers have the unique opportunity to support parents in their desire to spend time with their baby.
- Parents vary widely on how much time, if any, they want to spend with their baby, as well as when they'd like to see their baby, with whom, and what they would find meaningful and nurturing to do.
- Rather than having a single approach for all families or making sure that all bereaved parents spend time with their baby, the provider should engage in multiple conversations with the parents about their thoughts and feelings regarding their baby and the opportunity to see their baby, and then support them in doing what they believe is best.
- To support parents as they determine what to do and how to make the most of this irreplaceable time, the provider should offer unbiased information and individualized guidance, including exploring the options and possibilities, offering reassurances for their concerns, sharing what other parents have found helpful to do, and letting them know their baby continues to be freely available to them.
- Healthcare providers have a responsibility to make the baby freely available to parents for the duration of the hospital stay, as most parents are grappling with shock upon their baby's death and need plenty of time and support to consider their options. Some reluctant parents may change their minds, and knowing that their baby continues to be available gives them the freedom to reconsider.
- Whether parents want to be with their baby or not, parents need an experienced, knowledgeable, compassionate bereavement care provider to offer a safe and sacred space where they can do what is emotionally and spiritually meaningful to them.

Pregnancy Loss and Infant Death Alliance

SUPPORT . ADVOCACY . AWARENESS . EDUCATION

- Cultivating an empathic relationship with the parents is what enables the provider to ask them about their unique emotional, physical, cultural, and spiritual needs, and to be responsive to them.

Background and Significance

In the past few decades, increased awareness and sensitivity to the special needs of bereaved parents has changed hospital-based intervention—from *shielding* parents from the death of their child to *supporting* parents through the experience. As a result, the modern standard of care is to encourage and offer grieving parents repeated and extended opportunities to have close contact with their baby. Most parents value the time spent with their baby, as well as the photographs and keepsakes created during this time. Particularly for the mother, such contact can satisfy a strong and natural desire to provide ongoing care for the baby in whom she is emotionally invested and psychobiologically oriented.

However, this intervention is sometimes carried out insensitively or inconsistently, as some healthcare providers feel unsure, unsupported, or uncomfortable about accompanying parents through this experience. Some parents report being given little choice regarding their care, with a lack of communication and support from their providers. There have also been questions raised about the long-term benefit to parents, particularly as there are some parents who do not want to see their baby after death, and there is no published, empirical evidence that warrants insisting that reluctant parents do so. Finally, poor outcomes may be more likely for parents who do not receive adequate or appropriate follow-up care, underscoring the parents' need for information and long-term support around the grieving process as well as subsequent pregnancy and parenting. Understanding the complexities of parental bereavement requires further systematic research.

In the meantime, how can the healthcare provider navigate these tensions, and accommodate the parents who desire contact and stand to benefit *as well as* the parents who wish to decline time with their baby? And how can the provider work to ensure a better experience *and* outcome for all parents? The answer lies in focusing on process—that is, how the intervention is carried out. The process is supported by protocols, standard operating procedures, and policies with strong evidence supporting checklists as a framework for quality, consistent, and safe care practices.

Implications for Practice

Because many parents embrace the opportunity to spend time with their babies and report that they benefit from it, healthcare providers play a critical role in offering and supporting parents through this experience. However, offering the baby needs to be more than a direct or one-time query. Most parents are in shock, unfamiliar with the concept, and unable to weigh their needs on the spot. Instead, offering the baby should happen in the context of creating a caring relationship with the parents, in which the provider engages parents in periodic, open-ended conversations regarding

Pregnancy Loss and Infant Death Alliance

SUPPORT . ADVOCACY . AWARENESS . EDUCATION

their thoughts and feelings about their baby and about spending time with their baby's body. These conversations can help parents move through their shock, ask questions, get reassurance for their fears, consider their options, and identify their needs.

When parents want to see their baby, these conversations also allow the provider to help them determine how they want to spend this time. When one or both parents remain satisfied with their choice to *not* see their baby, the provider can support this decision and remain available to listen to parents talk about their baby and their grief. When a parent remains undecided, this indicates the need for more time and more dialogue. By remaining unbiased, openly exploring ideas and feelings, and letting the parents know their baby remains freely available, the provider can know that there is truly informed consent, whatever the parents decide to do.

Protocols and checklists, when used in the context of caring relationships, are important implements for promoting and guiding care. Protocols and checklists ensure choices and opportunities are offered to all parents regardless of the time of day, day of the week, or who is working. However, it is important that the healthcare provider must also individualize care, engaging parents in conversations about their options, enlisting each parent's right to make decisions, accommodating a variety of grief reactions, and regarding the uniqueness of each parent, family, need, and circumstance. Cultivating a caring relationship with parents is what makes this individualized care possible and effective.

In short, implementation of this intervention requires an experienced, knowledgeable, and compassionate bereavement care provider who sensitively engages parents around the opportunity to spend time with their baby, and then supports parents in proceeding at their own pace and in their own way. This focus on building relationships is known as relationship-based care, and is a way to ensure parent satisfaction with care during this tragic time.

Pregnancy Loss and Infant Death Alliance

SUPPORT . ADVOCACY . AWARENESS . EDUCATION

References

- Armstrong, D. S. (2007). Perinatal loss and parental distress after the birth of a healthy infant. *Advances in Neonatal Care*, 7(4), 200–206.
- Barnsteiner, J. H., & Gillis-Donovan, J. (1990). Being related and separate: A standard for therapeutic relationships. *MCN: The American Journal of Maternal Child Nursing*, 15(4), 223–228.
- Black, B. P., Wright, P. M., & Limbo, R. (Eds). (2016). *Perinatal and Pediatric Bereavement in Nursing and Other Health Professions*. New York, New York: Springer Publishing Company.
- Cacciatore, J. (2010). Stillbirth: Patient-centered psychosocial care. *Clinical Obstetrics & Gynecology*, 53(3), 691–699.
- Capitulo, K. L. (2005). Evidence for healing interventions with perinatal bereavement. *MCN: The American Journal of Maternal Child Nursing*, 30(6), 389–396.
- Carlson, R., Weber-Dennigmann, K., & Lammert, C. (2007). *Sharing and caring: Establishing and maintaining an effective Share program*. St. Charles, Missouri: Share Pregnancy and Infant Loss Support, Inc.
- Davis, D. L. (2016). *Empty cradle, broken heart: Surviving the death of your baby* (3rd ed.). Golden, Colorado: Fulcrum.
- Davis, D. L., & Helzer, S. (2010). Perinatal death and bereavement care. In E. S. Gilbert, (Ed.), *Manual of high-risk pregnancy & delivery* (5th ed.). New York, New York: Mosby/Elsevier.
- Davis, D. L., Stewart, M., & Harmon, R. J. (1988). Perinatal loss: Providing emotional support for bereaved parents. *Birth*, 15(4), 242–246.
- Gawande, A. (2010). *The checklist manifesto: How to get things right*. New York, New York: Picador.
- Gold, K. J. (2007). Navigating care after a baby dies: A systematic review of parent experiences with health providers. *Journal of Perinatology*, 27(4), 230–237.
- Gold, K., Dalton, V., & Schwenk, T. (2007). Hospital care for parents after perinatal death. *Obstetrics and Gynecology*, 109(5), 1156–1166.
- Health Service Executive, Bereavement Care Standards Development Group. (2016). National standards for bereavement care following pregnancy loss and perinatal death. Retrieved from <http://www.hse.ie/eng/about/Who/acute/bereavementcare/standardsBereavementCarePregnancyLoss.pdf>

Pregnancy Loss and Infant Death Alliance

SUPPORT . ADVOCACY . AWARENESS . EDUCATION

Hennegan, J. M., Henderson, J., & Redshaw, M. (2015). Contact with the baby following stillbirth and parental mental health and well-being: A systematic review. *BMJ Open*, 5, e008616.

Hughes, P., & Riches, S. Psychological aspects of perinatal loss. *Current Opinion in Obstetrics & Gynecology*, 15(2), 107–111.

Kavanaugh, K., & Hershberger, P. (2005). Perinatal loss in low-income African American parents. *Journal of Obstetric, Gynecologic, and Neonatal Nursing*, 34(5), 595–605.

Kavanaugh, K., & Moro, T. (2006). Supporting parents after stillbirth or newborn death: There is much that nurses can do. *The American Journal of Nursing*, 106(9), 74-79.

Kingdon, C., Givens, J. L., O'Donnell, E., & Turner, M. (2015). Seeing and holding baby: Systematic review of clinical management and parental outcomes after stillbirth. *Birth*, 42(3), 206-218.

Kobler, K., Limbo, R., & Kavanaugh, K. (2007). Meaningful moments: The use of ritual in pediatric and perinatal death. *MCN: The American Journal of Maternal Child Nursing*, 32(5), 288–296.

Koloroutis, M. (Ed.). (2004). *Relationship-based care: A model for transforming practice*. Minneapolis, Minnesota: Creative Health Care Management.

Koopmans, L., Wilson, T., Cacciatore, J., & Flenady, V. (2013). Support for mothers, fathers, and families after perinatal death. *The Cochrane Database of Systematic Reviews*, 19(6), CD000452.

Leon, I. G. (1992). Providing versus packaging support for bereaved parents after perinatal loss. *Birth*, 19(2), 89–91.

Limbo, R., & Kobler, K. (2010). The tie that binds: Relationships in perinatal bereavement. *MCN: The American Journal of Maternal Child Nursing*, 35(6), 316–321.

Limbo, R., Toce, S., & Peck, T. (2016). *Resolve Through Sharing® (RTS) position paper on perinatal palliative care* (Rev. Ed.). La Crosse, WI: Gundersen Lutheran Medical Foundation, Inc.

Limbo, R. K., & Wheeler, S. R. (1986/2003). *When a baby dies: A handbook for healing and helping* (Rev. ed.). La Crosse, Wisconsin: Gundersen Lutheran Medical Foundation, Inc.

Pregnancy Loss and Infant Death Alliance

SUPPORT . ADVOCACY . AWARENESS . EDUCATION

- Lisy, K., Peters, M. D., Riitano, D., Jordan, Z., & Aromataris, E. (2016). Provision of meaningful care at diagnosis, birth, and after stillbirth: A qualitative synthesis of parents' experiences. *Birth, 43*(1), 6-19.
- Meert, K. L, Thurston, C. S., & Briller, S. H. (2005). The spiritual needs of parents at the time of their child's death in the pediatric intensive care unit and during bereavement: A qualitative study. *Pediatric Critical Care Medicine, 6*(4), 420-427.
- O'Connell, O., Meaney, S., & O'Donoghue, K. (2016). Caring for parents at the time of stillbirth: How can we do better? *Women and Birth, pii: S1871-5192(16), 00006-00008*.
- O'Leary, J. Grief and its impact on prenatal attachment in the subsequent pregnancy. *Archives of Women's Mental Health, 7*(1), 7-18.
- Peters, M. D., Lisy, K., Riitano, D., Jordan, Z., & Aromataris, E. (2016). Providing meaningful care for families experiencing stillbirth: A meta-synthesis of qualitative evidence. *Journal of Perinatology, 36*(1), 3-9.
- Peters, M. D., Lisy, K., Riitano, D., Jordan, Z., & Aromataris, E. (2015). Caring for families experiencing stillbirth: Evidence-based guidance for maternity care providers. *Women and Birth, 28*(4), 272-278.
- Saflund, K., Sjogren, B., & Wredling, R. (2004). The role of caregivers after a stillbirth: Views and experiences of parents. *Birth, 31*(2), 132-137.
- Swanson, K. M. (1993). Nursing as informed caring for the well-being of others. *Image: The Journal of Nursing Scholarship, 25*(4), 352-357.
- Walling, A. D. (2002, November). *Tips from other journals: Should mothers see their infants after stillbirth?* Retrieved from American Family Physician website: <http://www.aafp.org/afp/2002/1115/p1956.html>
- Warland, J., & Davis, D. L. (2011). *Caring for families experiencing stillbirth: A unified position statement on contact with the baby. An international collaboration.* Retrieved from the MISS Foundation website: http://missfoundation.org/news/StillbirthContactwBaby_position_statement.pdf
- Workman, E. (2001). Guiding parents through the death of their infant. *Journal of Obstetric, Gynecologic, and Neonatal Nursing, 30*(6), 569-573.

Pregnancy Loss and Infant Death Alliance

SUPPORT . ADVOCACY . AWARENESS . EDUCATION

© 2008–2016 by Pregnancy Loss and Infant Death Alliance (PLIDA)

We invite you to use this position statement as a reference for articles, standard operating procedures, policies, and protocols. This document may be reprinted in its entirety without alterations. This document may be reprinted in its entirety without alterations. Verbatim portions of the statement or paraphrasing are permissible when a citation is included:

Adzich, K., Davis, D., Hochberg, T., Kavanaugh, K., Kobler, K., Lammert, C. A., . . . Press, J. N. (2016). *Pregnancy Loss and Infant Death Alliance (PLIDA) position statement on offering the baby to bereaved parents with relationship-base care* (Rev. ed.). Retrieved from the PLIDA website: <http://www.plida.org/position-statements/>

Contributors

Kathy Adzich, BA

Founder, Jakob's Room Project
www.trustingthejourney.com

Deborah L. Davis, PhD

Developmental Psychologist and Writer
Denver, Colorado

Todd Hochberg, BA

Bereavement Photographer
Touching Souls Photography
www.touchingsouls.org

Karen Kavanaugh, PhD, RN, FAAN

Elizabeth Schotanus Professor of Pediatric Nursing
Wayne State University

Kathie Kobler, MS, APN, PCNS-BC, CHPPN, FPCN

Pediatric Palliative & Supportive Care
Center for Fetal Care
Advocate Children's Hospital

Catherine A. Lammert, RN

Education Coordinator (retired)
Share Pregnancy and Infant Loss Support, Inc.

Irving Leon, PhD

Clinical Psychologist
Adjunct Associate Professor of Obstetrics and Gynecology
University of Michigan Health System

Rana Limbo, PhD, RN, CPLC, FAAN

Associate Director and Senior Faculty Consultant
Resolve Through Sharing®
Bereavement and Advance Care Planning Services
Gundersen Medical Foundation, Inc.

Darryl Owens, MDiv, BCC, CT

Women's Services Chaplain/Grief Counselor
Clinical Chaplain II, University of North Carolina Hospitals

Janet N. Press, CNS, MSN, CT, RN-C

Perinatal Obstetrical Coordinator
Central New York Region
Perinatal Bereavement Services Coordinator
Crouse Hospital